

Is birthing pain the trigger of postpartum depression?

Pascal H. Vuilleumier, MD^a, Sarah W. Prager, MD^b

a) University Department of Anesthesiology and Pain Therapy, Bern University Hospital, Inselspital, Bern, Switzerland

b) Department of Obstetrics and Gynecology,
University of Washington Medical Center, Seattle, USA

Corresponding address:

Pascal H. Vuilleumier, M.D.

Department of Anesthesiology and Pain Medicine

Bern University Hospital

Inselspital

3010 Bern, Switzerland

Phone: +41 31 632 17 09, Fax: +41 31 632 05 54

Email: pascal.vuilleumier@insel.ch

Keywords: childbirth, pain, postpartum depression, pain management

1

2 **Background**

3

4 In 2007 the International Association for the Study of Pain (IASP) launched the
5 campaign "Real Women, Real Pain"; 2007-2008 was declared as the "Global Year
6 Against Pain in Women. By then it was recognized that 90% of women experience
7 severe/unbearable labor pain, and that one very real consequence to labor pain was
8 psychological: "Long-term emotional stress with potential adverse consequences on
9 maternal mental health and family relationships" [1]. One year later Eisenach and
10 his colleagues published their seminal data on the association between acute pain
11 and postpartum depression [2]. Eisenach conducted a multicentre study across
12 North America and Western Europe, recruiting 1288 women after vaginal and
13 cesarean delivery (CD). The reported combined incidence of acute pain 36 hours
14 after delivery was 10.6%, and persistent pain at 8 weeks was 9.8%. Multivariable
15 regression revealed a threefold increased risk of postpartum depression (PPD) in
16 the presence of acute pain.

17 Rates of persistent pain after cesarean delivery (PPCD) is subject to debate [3]. On
18 the lower end, Eisenach et al. reports PPCD as 1.8% at 6 months decreasing to 0.3% at 12
19 months [4]. On the other end of the spectrum, 20% of PPCD is reported 2 years after CD
20 for breech lie [5]. Other studies report rates between 6-18% beyond 6 months postpartum
21 [6-8]. Although vaginal birth is one of the most frequent significant life events in women,
22 persistent pain after vaginal delivery (PPVD) is more difficult to quantify and correlate

1 across studies than PPCD, resulting in more limited data. One study reports an incidence
2 of disabling PPVD in 2-5% of women two years after delivery [9]; another reports an
3 incidence of 1% PPVD at one year [10]. Although the degree of pelvic trauma or
4 episiotomy does not appear to correlate with PPVD at 6 weeks [3, 11, 12], the
5 experience of assisted vaginal birth (vacuum extraction or forceps) seems to
6 increase PPVD rates [5, 7, 13].

7 The prevalence of some degree of emotional disturbances in parturient women has
8 been cited to be as high as 85% [14]. This almost equals the incidence of severe or
9 unbearable pain during the birth process, which has been cited to be 90% [1]. The
10 DSM-5 now categorizes PPD as a diagnosed depressive disorder with peripartum
11 onset (within four weeks after delivery) [15]. The diagnosis of PPD still lacks a clear
12 definition [16, 17], though it's prevalence is typically cited to be between 10-20%
13 [18]. In western countries, most mothers leave the hospital 24-72 hours after
14 delivery, typically before PPD would manifest. Most PPD is diagnosed in an
15 outpatient setting, resulting in under-diagnosis of PPD. One study including 43,093
16 postpartum women has reinforced the rates of postpartum psychiatric illness, but
17 also revealed low overall rates of pregnant and postpartum women seeking and/or
18 receiving care for chronic or new mental health [19].

1 Discussion

2
3 Today only three prospective and one retrospective studies were able to detect a
4 significant correlation between pain during the childbirth and PPD (table 1).

5 The study by Hiltunen et al was a prospective follow-up comparing 162 mothers
6 during the first postpartum week and at four months [20]. This study shows a
7 positive correlation between analgesia and PPD one week after childbearing, but the
8 study lacks a measured pain outcome. Additionally, the analysis differentiates by
9 type of anesthesia provided, but doesn't account for quality of analgesia,
10 constituting another serious limitation to this study.

11 The primary aim of Eisenach's prospective study was to explore the consequences
12 of cesarean delivery on persistent pain at 8 weeks. This study revealed that the
13 perceived severity of acute pain, but not the mode of delivery, was a significant risk
14 factor for both persistent pain and postpartum depression [21]. Postpartum
15 depression was actually a secondary outcome in this study. This may actually be a
16 strength, since the focus on pain may help eliminate some confounding bias that has
17 limited other studies.

18 The retrospective study by Gaudet et al is the largest dataset to date, correlating
19 postpartum pain at 3 and 7.3 months to PPD (OR 1.7, 95% CI 1.2-1/5; OR 2.4, 95%
20 CI 1.6-3.6, respectively) [22]. Again, this study was limited by lack of a precise pain
21 measurement and no discrimination between types of analgesia offered during
22 labor and delivery.

1 The most complete prospective study was conducted by Ding et al in Beijing,
2 revealing an OR of 0.31 for PPD in subjects who received epidural analgesia during
3 labor [21]. Epidural analgesia was the only available form of pain management
4 during labor, so that was standardized and the study population was dichotomized
5 between epidural and "no pain relief at all" groups. The statistical analysis
6 accounted for significant confounding factors, including breastfeeding, Edinburgh
7 Postnatal Depression Scale (EPDS) scores at 3 days postpartum and maximal pain
8 scores during complete cervical dilation.

9 These four studies examine the association between PPD and severe pain during
10 childbirth. Psychological and environmental factors for postpartum depression have
11 been well described [17]. More recently, there is an acknowledgement by Wisner
12 [23] who, in commenting on the study by Ding [21], reveals a possible psychological
13 element also to biological factors that may be related to a link between PPD and
14 childbirth pain.

15 One biological cause of PPD in susceptible women that is an area of intense research
16 interest is the rapidly fluctuating levels of estrogen and progestogen occurring at
17 birth [24]. Alternatively, exposure to oxytocin/oxytocin agonists during and after
18 labor may reduce PPD. Gutierrez et al have shown in a rodent model that oxytocin
19 has an analgesic effect during the puerperium [25]. It is now well established that
20 oxytocin-induced nociception is mediated by a subpopulation of glutamatergic
21 neurons amplifying GABAergic inhibition of pain [26]. One major drawback with
22 oxytocin studies is that oxytocin does not cross the blood-brain barrier, but another
23 oxytocin agonist, carbetocin, has been shown to have analgesic effects when

1 compared to oxytocin administration after cesarean delivery [27]. Recently
2 intravenous carbetocin has shown anti-hyperalgesic effects in healthy volunteers
3 [28]. All of these potential biologic links between peripartum pain and PPD seem to
4 be influenced by the individual's perception and experience of pain, not just
5 presence or degree of pain. This would certainly help account for cultural
6 differences in management of pain and prevalence of PPD [28].
7
8

9 **Conclusion**

10
11 In the last 20 years obstetric anesthesia and analgesia has made a significant leap
12 forward in understanding and treating pain during the birth process. Techniques for
13 either epidural analgesia used during labor or neuraxial anesthesia administered for
14 cesarean delivery have dramatically progressed, paralleled by a gain in popularity.
15 20 years ago the standard for pain management during labor was administration of
16 nitrous oxide and general anesthesia for cesarean delivery. Today we understand
17 that postpartum depression and pain during labor are two complex, multifactorial,
18 and, to some degree, interrelated problems. Individually, each is considered a major
19 health issue and investigators are now looking more closely on how peripartum
20 pain and PPD are associated. Oxytocin, sometimes known as the "love hormone", is
21 known to be an important promoter of mother-child bonding; perhaps it will also
22 prove to be clinically useful in preventing or treating postpartum depression [29].

1 Although further work is needed in order to understand how pain and depression
2 relate to each other, recent publications suggest that optimal desired pain control
3 during the birth process may decrease the prevalence of postpartum depression.
4 Improved attention to helping women achieve desired pain control during and after
5 childbirth may have far-reaching benefits.

6

7

8

1

2 **References**

3

4 1. Charlton JE, editor. Pain and Pregnancy and Labor. 3rd Edition ed. Seattle:
5 IASP Press; 2005.

6 2. Eisenach JC, Pan PH, Smiley R, Lavand'homme P, Landau R, Houle TT.
7 Severity of acute pain after childbirth, but not type of delivery, predicts persistent
8 pain and postpartum depression. Pain. 2008;140(1):87-94. Epub 2008/09/27. doi:
9 10.1016/j.pain.2008.07.011. PubMed PMID: 18818022; PubMed Central PMCID:
10 PMC2605246.

11 3. Lavand'homme P. Chronic pain after childbirth. Current Opinion in
12 Anaesthesiology. 2013;26(3):273-7. doi: 10.1097/ACO.0b013e328360c57b.
13 PubMed PMID: 23549199.

14 4. Eisenach JC, Pan P, Smiley RM, Lavand'homme P, Landau R, Houle TT.
15 Resolution of pain after childbirth. Anesthesiology. 2013;118(1):143-51. doi:
16 10.1097/ALN.0b013e318278ccfd. PubMed PMID: 23249931.

17 5. Hannah ME, Whyte H, Hannah WJ, Hewson S, Amankwah K, Cheng M, et al.
18 Maternal outcomes at 2 years after planned cesarean section versus planned vaginal
19 birth for breech presentation at term: The International Randomized Term Breech
20 Trial. American Journal of Obstetrics and Gynecology. 2004;191(3):917-27. doi:
21 10.1016/j.ajog.2004.08.004. PubMed PMID: 15467565.

- 1 6. Nikolajsen L, Sorensen HC, Jensen TS, Kehlet H. Chronic pain following
2 Caesarean section. *Acta Anaesthesiologica Scandinavica*. 2004;48(1):111-6. PubMed
3 PMID: 14674981.
- 4 7. Declercq E, Cunningham DK, Johnson C, Sakala C. Mothers' reports of
5 postpartum pain associated with vaginal and cesarean deliveries: results of a
6 national survey. *Birth*. 2008;35(1):16-24. doi: 10.1111/j.1523-536X.2007.00207.x.
7 PubMed PMID: 18307483.
- 8 8. Kainu JP, Sarvela J, Tiippana E, Halmesmaki E, Korttila KT. Persistent pain
9 after caesarean section and vaginal birth: a cohort study. *International Journal of*
10 *Obstetric Anesthesia*. 2010;19(1):4-9. doi: 10.1016/j.ijoa.2009.03.013. PubMed
11 PMID: 19733050.
- 12 9. Albert H, Godskesen M, Westergaard J. Prognosis in four syndromes of
13 pregnancy-related pelvic pain. *Acta Obstet Gynecol Scand*. 2001;80(6):505-10.
14 PubMed PMID: 11380285.
- 15 10. Eisenach JC, Pan P, Smiley RM, Lavand'homme P, Landau R, Houle TT.
16 Resolution of pain after childbirth. *Anesthesiology*. 2013;118(1).
- 17 11. Carroli G, Mignini L. Episiotomy for vaginal birth. *The Cochrane Database of*
18 *Systematic Reviews*. 2009;(1):CD000081. doi: 10.1002/14651858.CD000081.pub2.
19 PubMed PMID: 19160176; PubMed Central PMCID: PMC4175536.
- 20 12. Macarthur AJ, Macarthur C. Incidence, severity, and determinants of perineal
21 pain after vaginal delivery: a prospective cohort study. *American Journal of*
22 *Obstetrics and Gynecology*. 2004;191(4):1199-204. doi:
23 10.1016/j.ajog.2004.02.064. PubMed PMID: 15507941.

- 1 13. Thompson JF, Roberts CL, Currie M, Ellwood DA. Prevalence and persistence
2 of health problems after childbirth: associations with parity and method of birth.
3 Birth. 2002;29(2):83-94. PubMed PMID: 12051189.
- 4 14. Cotichelli G, Barbadoro P, Chiatti C, Marigliano A, Prospero E, Di Stanislao F.
5 Regional distribution of postpartum depressive symptoms in Italy. The 15th
6 Emerging New Researchers in the Geography of Health and Impairment Conference;
7 June 10-11 2010; Paris 2010.
- 8 15. Association AP. Diagnostic Manual of Mental Disorders. Fifth ed. ed:
9 American Psychiatric Publishing; 2013.
- 10 16. Wisner KL, Moses-Kolko EL, Sit DK. Postpartum depression: a disorder in
11 search of a definition. Arch Womens Ment Health. 2010;13(1):37-40. doi:
12 10.1007/s00737-009-0119-9. PubMed PMID: 20127453; PubMed Central PMCID:
13 PMC4426488.
- 14 17. O'Hara MW, Wisner KL. Perinatal mental illness: definition, description and
15 aetiology. Best Practice & Research Clinical Obstetrics & Gynaecology. 2014;28(1):3-
16 12. doi: 10.1016/j.bpobgyn.2013.09.002. PubMed PMID: 24140480.
- 17 18. Miller LJ. Postpartum depression. JAMA 2002;287(6):762-5. PubMed PMID:
18 11851544.
- 19 19. Vesga-Lopez O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric
20 disorders in pregnant and postpartum women in the United States. Archives of
21 General Psychiatry. 2008;65(7):805-15. doi: 10.1001/archpsyc.65.7.805. PubMed
22 PMID: 18606953; PubMed Central PMCID: PMC2669282.

- 1 20. Hiltunen P, Raudaskoski T, Ebeling H, Moilanen I. Does pain relief during
2 delivery decrease the risk of postnatal depression? *Acta Obstet Gynecol Scand*.
3 2004;83(3):257-61. Epub 2004/03/05. doi: 302 [pii]. PubMed PMID: 14995921.
- 4 21. Ding T, Wang DX, Qu Y, Chen Q, Zhu SN. Epidural labor analgesia is associated
5 with a decreased risk of postpartum depression: a prospective cohort study.
6 *Anesthesia and Analgesia*. 2014;119(2):383-92. Epub 2014/05/07. doi:
7 10.1213/ANE.0000000000000107. PubMed PMID: 24797120.
- 8 22. Gaudet C, Wen SW, Walker MC. Chronic perinatal pain as a risk factor for
9 postpartum depression symptoms in Canadian women. *Can J Public Health*.
10 2013;104(5):e375-87. Epub 2013/11/05. PubMed PMID: 24183178.
- 11 23. Wisner KL, Stika CS, Clark CT. Double duty: does epidural labor analgesia
12 reduce both pain and postpartum depression? *Anesthesia and Analgesia*.
13 2014;119(2):219-21. Epub 2014/07/22. doi: 10.1213/ANE.0000000000000322
14 00000539-201408000-00001 [pii]. PubMed PMID: 25046776.
- 15 24. Pearlstein T, Howard M, Salisbury A, Zlotnick C. Postpartum depression.
16 *American Journal of Obstetrics and Gynecology*. 2009;200(4):357-64. doi:
17 10.1016/j.ajog.2008.11.033. PubMed PMID: 19318144; PubMed Central PMCID:
18 PMC3918890.
- 19 25. Gutierrez S, Liu B, Hayashida K, Houle TT, Eisenach JC. Reversal of peripheral
20 nerve injury-induced hypersensitivity in the postpartum period: role of spinal
21 oxytocin. *Anesthesiology*. 2013;118(1):152-9. doi:
22 10.1097/ALN.0b013e318278cd21. PubMed PMID: 23249932; PubMed Central
23 PMCID: PMC3876486.

- 1 26. Breton JD, Veinante P, Uhl-Bronner S, Vergnano AM, Freund-Mercier MJ,
2 Schlichter R, et al. Oxytocin-induced antinociception in the spinal cord is mediated
3 by a subpopulation of glutamatergic neurons in lamina I-II which amplify GABAergic
4 inhibition. *Molecular Pain*. 2008;4:19. Epub 2008/05/31. doi: 10.1186/1744-8069-
5 4-19. PubMed PMID: 18510735; PubMed Central PMCID: PMC2430948.
- 6 27. De Bonis M, Torricelli M, Leoni L, Berti P, Ciani V, Puzzutiello R, et al.
7 Carbetocin versus oxytocin after caesarean section: similar efficacy but reduced
8 pain perception in women with high risk of postpartum haemorrhage. *The Journal*
9 *of Maternal-fetal & Neonatal medicine*. 2012;25(6):732-5. doi:
10 10.3109/14767058.2011.587920. PubMed PMID: 21761999.
- 11 28. Vuilleumier PH, Schliessbach J, Biurrun Manresa JA, Müller M, Stamer UM,
12 Arendt-Nielsen L, et al. Analgesic effects of oxytocin receptor modulation in healthy
13 volunteers. 2015 EFIC Congress: Pain in Europe IX; 9.2-5 2015; Vienna2015.
- 14 29. Zik JB, Roberts DL. The many faces of oxytocin: implications for psychiatry.
15 *Psychiatry Res*. 2015;226(1):31-7. doi: 10.1016/j.psychres.2014.11.048. PubMed
16 PMID: 25619431.

17

Table 1: Overview of studies significantly correlating postpartum pain to postpartum depression

Year	Correlation of pain and postpartum depression	N	Groups compared	PPD	Acute pain during birth	PPD	Reference
2004	OR 0.25, (95%CI 0.09-0.72) for PPD at one week in analgesia group compared with no analgesia	162	-No analgesia (N=23) -Nitrous oxide or acupuncture (N=16) -Epidural or paracervical block (N=103) -Elective cesarean (N=32) -Emergency cesarean (N=11)	EPDS \geq 13	NA	13% at 4 months	[20]
2008	Every NRS point increase in acute pain is correlated to a 8.3% increase in 8-week EDPS p<0.001	1288	-Vaginal delivery (N=837) -Cesarean delivery (N=391)	EPDS>12	NRS 3.3 (SD 2.1) NRS 4.7 (SD 2.0)	11.2% at 2 months	[2]
2013	OR 1.7, (95%CI 1.2-1.5) for PPD if perinatal pain present at 3 months, OR 2.4, 95%CI 1.6-3.6 for PPD if problematic perineal pain at 7.3 months	5614	-Presence of problematic perinatal pain in the first 3 months -Duration of problematic perinatal pain -Number of types of perinatal pain at interview	EPDS \geq 13	81.7% at 3 months	7% at 7.3months	[22]
2014	OR 0.31, 95%CI 0.12-0.82) for PPD in epidural group	214	-Labor epidural (N=107) -No labor epidural (N=107)	EPDS \geq 10	NRS 3 (SD 0-7) NRS 10 (SD 7-10)	24.3% at 6 weeks	[21]

Table legend: OR= Odds ratio, CI=Confidence interval, PPD=Postpartum depression, N=Number cases, EPDS=Edinburgh postpartum depression scale, NRS=Numeric rating scale of pain, SD=Standard deviation, NA=Not available